Resistance in Psychotherapy

Allison Felton, MD
Assistant Professor, University of Missouri-Columbia School of Medicine

Sources of Resistance

- Patient originated
- Therapist originated
- Social network originated

Patient Variables

- Readiness to Change
- Lack of Information
- Fears associated with change
- Dysfunctional patterns expressing themselves during therapy
- Transference

Patient Originated Resistance

 The Usual Response to fear is what?

Psychotherapy/ Pharmachotherapy

- Patient unsure of taking a medication
- Recognition of pattern of fear when patient taking on anything new
- Interpretation of patients fears in regards to medication
- Recognition of patterns inv

Patient Originated

- Lack of insight-can't identify feelings or recognize patterns
- Dysfunctional patterns (fear of criticism, not doing homework perfectly)
- The pain of where they are is NOT greater than the pain of change

Therapist Variables

- Inappropriate expectations of patient
- Lack of skills or lack of skills that fit with patient
- Poor explanation of homework
- Working toward the wrong goals
- Countertransference

Social Network Variables

- Criticism of patient being in therapy
- Upsetting the equilibrium of the social network
- When one member of a unit changes others are changed whether they like it or not

Recognition of Resistance-Therapist

Frustration or discomfort

Expectation of Resistance

 Need to understand various forms of resistance in relationship to stage of psychotherpay

Engagement Stage- Forms of Resistance

- Difficulty with trusting the therapist
- Doubting the effectiveness of therapy
- DNKA
- Patient criticizing therapist

Pattern Search Stage-Forms of Resistance

- Failure to report information relevant to defining patterns as seen by:
- -silence
- -defensiveness
- -lying
- -not doing homework
- -talking about irrelevant topics

Homework

Change stage-Forms of Resistance

- Failure to generate alternatives as evidenced by:
- -agree to try to generate new pattern but fail to do so
- -refusal to comply with therapists suggestions
- -discuss termination before changing

Termination Stage-Forms of Resistance

- Fail to terminate therapy such as:
- -continuing to call therapist
- -fear of loss of therapist means return of symptoms

Confronting Resistance-Therapist's Role

- Identification of resistance
- Contemplate origins
- Is this an example of a maladaptive pattern?
- Why am I so frustrated with this?
- What is motivating this person to respond in a self defeating way?
- Am I expecting too much from this patient?
- Did I explain exercises thoroughly?
- Does patient understand?

Confronting Resistance

- Therapist's need to look at their own expectations of the patient keeping in mind:
- 1. patient's readiness to change,
- 2. their degree of fear
- 3. social network support,
- 4. patient's functioning level

Beginning Therapists

- Use of poor interventions
- Knowledge base
- Unrealistic expectations

Social Network

- Friends drink-have to find new social interests or friends
- Woman develops chronic illnessfather of children resistant to taking more responsibility
- Husband stops drinking-wife can no longer blame his drinking for her lack of desire for intimacy

Stages of Psychotherapy